

Effect of Environmental Factors on Chemical Composition of Seeds in Soybean, Sesame, Linseed and Safflower Crops

Hafiz Saad Bin Mustafa^{*1}, Tariq Mahmood¹, Muhammad Aftab¹, Abdul Qayyum¹ and Muhammad Aslam²

¹Directorate of Oilseeds, Ayub Agricultural Research Institute (AARI), Faisalabad, Pakistan.

²Department of Plant Breeding & Genetics, University of Agriculture Faisalabad, Pakistan.

Corresponding author: saadpbg@gmail.com

Abstract

The aim of present review article is to clarify the effect of climate on seed chemical composition of Soybean, Sesame, Linseed and Safflower. Environment plays an important role during the growing stages of crops and eventually influence the chemical composition of seed. Temperature, rainfall, sowing time, irrigation, fertilizers, growing season and soil type during different growth stages affect chemical composition of different oilseed crops. Oil content and protein content are specially influenced by prevailing environmental conditions during maturity and post flowering period. High temperature during seed filling stage reduces oil content and increases protein content in safflower. Genotype and location has significant effect on protein, oil and unsaturated fatty acids, namely oleic acid, linoleic acid and linolenic acid in soybean. Oil content and oleic acid increase while protein, linoleic and linolenic acids decrease in early sown soybean. Due to drought stress, protein content increases while oil content decreases in soybean. Pod position also affects protein, oil and iso-flavon content of soybean. Oil content and composition in sesame is influenced by soil type, genetic, climatic and agronomic factors as well as growth stages of plant. Temperature variances after flowering express huge impact on content and composition of the linseed oil. Late sowing decreases oil content in linseed while cool temperature, adequate moisture and long day length during and after flowering increase the seed yield, quality and quantity of oil content. Drought stress at flowering and grain filling stages reduces oil content and un-saturated fatty acids, while increases saturated fatty acids and protein in safflower.

Key words: Drought, Genetic variation, Oil quality, Oilseed crops, Rainfall, Temperature

Introduction:

Pakistan is deficient in its edible oil requirement and spends huge amount annually on the import of edible oil, which badly affect the economy of the country. During the year 2016-17, the total availability of edible oil was 3.263 million tonnes. Local production of edible oil remained 0.431 million tonnes while import of edible oil or oilseeds was 3.191 million tonnes. The import bill during 2016-17 was Rs. 320.893 billion (US\$ 3.063 billion). *Economic Survey of Pakistan, 2017-18*.

Environmental conditions influence crop growth and development and are the most important yield reducing factors in the world (Franklin *et al.*, 2010). Environmental factors during flowering and seed-filling period affect seed yield and seed quality of oilseed crops (Ali *et al.*, 2009; Monotti, 2003; Petcu *et al.*, 2001). Genotypes behave differently under different environmental conditions (Qadir *et al.*, 2006). Fatty acid composition of oil varies according to the environmental conditions (Strecker *et al.*, 1997; Pritchard *et al.*, 2000; Roche *et al.*, 2006). Accumulation of saturated fatty acids is also affected by environmental factors (Roche *et al.*, 2006; Izquierdo and Aguirrezábal, 2008). The effect of temperature on the fatty acid composition of the seed oil has been studied in different oilseed crops, including flax (Yermanos and Gooden, 1965; Green,

1986) and soybean (Martin *et al.*, 1986). Ecological factors other than temperature, such as solar radiation (Santalla *et al.*, 1995), precipitation (Pritchard *et al.*, 2000), nitrogen availability (Steer and Seiler, 1990), soil salinity (Irving *et al.*, 1988) and crop vigor (Zimmer and Zimmerman, 1972), can also affect fatty acid composition. Post-flowering temperatures have a major effect on fatty acid profile of different oilseed crops (Pritchard *et al.*, 1999). The fatty acid composition in different oilseed crops is influenced by the temperature prevailing during seed maturation (Hilditch and Williams, 1964). Inconsistent ecological conditions may result in wide variation in oil quantity and composition in different oilseeds (Shafii *et al.*, 1992). Prevalence of lower temperature at the time of maturation resulted in an increase in polyunsaturated fatty acids in oilseed plants (Nytker *et al.*, 2006). Each 1°C increase of temperature leads to about 2% increase of oleic acid (Demurin *et al.*, 2000).

Temperature is an important environmental factor affecting the fatty acid composition of different plant portions like seeds (Tremolieres *et al.*, 1978), roots (Simolenka and Kuiper, 1977) and leaves (Wilson and Crawford, 1974). Temperature differential has strong influence on proportions of different fatty acids (Matsuzaki *et al.*, 1988) and

molar proportion of oleic acid increased and that of linoleic and linolenic acid decreased as temperature increased. Tocopherol content in the oil of soybean (Izquierdo, 2007) depicted inverse relationship with the quantity of sun rays intercepted per plant during grain filling.

Climatic conditions, water and nitrogen availability mainly during the seed-filling stage account for most of the variations than due to genotype differences in grain protein concentration (Cooper *et al.*, 2001). Oil and protein are two major components of seeds in oilseed crops and are affected significantly by environmental stresses (Dehnavi and Sanavy, 2008). Up to 17% decrease in yields of soybean for 1°C increase in average growing season temperature above the optimum is reported in the United States (Lobell and Asner, 2003). Large amounts of oil are stored mainly in embryonic tissues (cotyledons) that occupy most of the space available in between the integuments of the seed in oilseeds crops including soybean, linseed and safflower (Baud and Lepiniec, 2010).

Soybean, Sesame, Linseed and Safflower are the minor oilseed crops of the country and contribute an important portion in the local oilseed production of Pakistan. Therefore, it is need to study the effect of climatic factors on seed chemical composition of these crops.

SOYBEAN

Soybean (*Glycine max* L.) is important oil crop with 18-22% oil and 35-40% protein. It is known as the main source of plant oil and protein with highest acreage in the world (Maleki *et al.*, 2013). About 90% of the world's soybean production occurs under rainfed conditions characterized by high temperature and low or erratic rainfall (Thuzar *et al.*, 2010). Roughly 60% of soybean dry weight is composed of protein and oil, ranking it 1st for protein content and 2nd for oil content among other cereal and legume species (Liu, 1997). Soybean oil averages 12% palmitic acid (16:0), 4% stearic acid (18:0), 23% oleic acid (18:1), 53% linoleic acid (18:2), and 8% linolenic acid (18:3). The 16:0 and 18:0 fractions are saturated fatty acids and constitute 15% of the soybean oil. The remainder of the oil (about 85%) is made up of unsaturated fatty acids or 18:1, 18:2, and 18:3 (Lee *et al.*, 2007). Seed protein content is important for both feed and food utilization of soybean (Vollmann *et al.*, 2000).

Growing season significantly influence protein content in soybean (Vollmann *et al.*, 2000). Genetic and environmental factors are major determinants of grain yield, protein and oil concentration of soybean (Wolf *et al.*, 1982, Maestri *et al.*, 1998). Intercepted radiation showed no impact on grain oil concentration in soybean (Andrade and Ferreiro, 1996). Average to higher protein content was recorded in years with high air temperature and

moderate rates of rainfall during the grain filling period whereas seed protein content was drastically reduced (range 265±347 g kg⁻¹) in seasons of insufficient nitrogen fixation or higher amounts of precipitation during grain filling (Vollmann *et al.*, 2000). The proportion of large seeds was higher at the apical portion while small seeds predominated at the basal portion of the soybean stem axis. The contents of lipids, starch, soluble sugars and soluble proteins were higher in large seeds as compared to those in smaller ones. The percentage of membrane lipid components, on a 10-kernel basis, was higher in large seeds (Guleria *et al.*, 2008).

Seed oil (16%) and oleic acid (22.8%) increase while protein (6.6%), linoleic (10.9%) and linolenic acids (27.7%) decrease in early sown soybean than late planting under irrigated conditions (Bellaloui *et al.*, 2011). Late planting resulted in higher sucrose and raffinose and lower stachyose compared with early planting (Bellaloui *et al.*, 2011). Both, irrigation and planting date have a significant influence on seed protein, oil, unsaturated fatty acids and sugars (Bellaloui *et al.*, 2011). Oil concentration increased with early planting (Pedersen and Lauer, 2004; Kane *et al.*, 1997; Helms *et al.*, 1990). Protein concentration increased and oil concentration decreased with late planting (Kane *et al.*, 1997; Helms *et al.*, 1990). Palmitic and linolenic acids may decrease with later planting date, but stearic acid may increase and this may be due to temperature changes during seed maturation at later planting (Wilcox and Cavins, 1992). Oleic acid levels increased while those of linoleic and linolenic acid decreased when soybeans were grown in warmer environments (Carver *et al.*, 1986). Temperature may affect oleate and linoleate desaturases (Burton, 1991), decrease oleyl and linoleyl desaturase activities at 35°C (Cheesbrough, 1989), decrease ω-6 desaturase enzyme, encoded by the FAD2-1A gene, and desaturases degraded at high growth temperatures of 30°C (Tang *et al.*, 2005).

Seeds collected from the base part of soybean plants had greater pod length and pod width than those collected from the top part of plants (Al-Tawaha, 2010). Environmental conditions can influence the development of the seed and affect its composition (Cook, 2008). Adverse climatic conditions such as low temperature and high amounts of precipitation prevailing in northern regions of Europe reduce protein content (Vollmann *et al.*, 2000). In the USA, seed protein content in the Western and Eastern Corn Belt was clearly lower than in southern regions of production over a number of seasons (Hurburgh, 1994). Nian *et al.*, (1996) reported low protein content in the northern locations of north-east China and in northern sites of Europe, due to large seasonal variations (Schuster and Boehm 1981). Protein content showed significant negative ($P < 0.01$) while

oil content ($P < 0.05$) exhibited positive correlation with latitude (Kumar *et al.*, 2006). Vollman *et al.*, (2000) showed that during years of low precipitation and high temperature, protein synthesis was higher than oil synthesis in soybean seeds. Soil moisture level affects protein while the timing with respect to maturity of irrigation controls oil content (Foroud *et al.*, 1993). High rainfall during the grain filling stage can lead to higher oil content in soybean (Vollman 2000). Sucrose and stachyose level were affected by the environment and content of both decreased with increasing temperature in soybeans, with sucrose showing the steepest level of decline per degree increase (Wolf *et al.*, 1982).

Protein and oil content of soybean is determined by genetic and environmental factors and the stage of maturity of the grain (Wolf *et al.*, 1982; Maestri *et al.*, 1998; Baydar and Sabri, 2005). Composition of soybean seed varies with environmental factors prevailing during the seed filling period when accumulation of the seed chemical components occurs (Wolf *et al.*, 1982; Wilson, 2004; Carrera *et al.*, 2009, 2011). The relationship of protein contents with temperature (Piper & Boote, 1999; Wilson, 2004) and water availability (Rose, 1988; Boydak *et al.*, 2002; Kumar *et al.*, 2006; Carrera *et al.*, 2009) has been studied. Higher temperature favors the accumulation of sulphur amino acids, methionine and cystine (Wolf *et al.*, 1982). Soybean from Northern zones of the United States are cooler than central and southern zones showed lower essential, nonessential and total amino acid content (Grieshop and Fahey 2001 and Karr-Lilienthal *et al.*, 2005). Application of nitrogen fertilizers at various growth stages has proven ineffective in improving the protein or oil concentration of soybean (Singh *et al.*, 2001; Schmitt *et al.*, 2001; Wesley *et al.*, 1998). Rainfall showed a negative correlation with protein content. Average daily mean temperatures during bean development showed a positive correlation with protein and a negative one with oil and linolenic acid. Genotypic, locational and genotypic-locational interactions were found to be significant for protein, oil and unsaturated fatty acids, namely oleic acid, linoleic acid and linolenic acid (Kumar *et al.*, 2006). Increasing water stress at reproductive stages resulted in an increment in protein content but oil percentage decreased. Seeds present at upper position of the canopies showed higher oil and protein content than those from middle and lower parts under all irrigation treatments (Ghassemi-Golezani and Lotfi, 2013). Water deficiency has little effect on seed protein content in soybean (Thompson 1978) while Hobbs and Muendel (1983) reported that drought stress increases protein content in soybean. At reproduction stage drought stress results in decreased grain size in soybean (Momen *et al.*, 1979; Kadhem *et al.*, 1985). Bravedan and Egli (2003) reported significant decrease (39%) in

yield due to fewer and smaller grains if short period of water stress occurs during grain filling of soybean.

Seed position on mother plant is an important factor influencing grain quality (Ghassemi-Golezani and Lotfi, 2013). It may result in differences within-plant and account for part of the variation in physical (weight, shape) or physiological (germination and vigor) seed attributes (Illipronti 2000). Variation in protein content is reported among different nodes of same plant and protein content was lowest in the basal node grains and increases toward the apical nodes in soybean plants (Escalante and Wilcox 1993). Seed protein, oil quality and quantity of soybean showed significant variation with node position (Collins and Carter, 1956; Escalante and Wilcox, 1993). Indeterminate cultivars depicted more oil content in seeds from lower nodes than seeds on the upper nodes, while in the case of determinate cultivars oil content was highest in seeds from terminal nodes (Collins and Cartter, 1956). Seed from the middle nodes on determinate cultivars had higher oil and lower protein than seeds from either the top or bottom of the plants (Collins and Cartter, 1956). Lowest nodes in both determinate and indeterminate types of soybean showed lowest seed protein concentration. Linear increase in protein concentration from 397 g kg⁻¹ at the lowest node to 442 g kg⁻¹ at the highest node in determinate and increased from the lowest (398 g kg⁻¹) and increased progressively through 14 in indeterminate, with no significant differences between nodes 12 and 16 (Escalante and Wilcox, 1993). Concentration of S in soybean leaves decreases during grain filling (Sweeney and Granade, 1993, Sexton *et al.*, 1998) as does N/S ratio (Sweeney and Granade, 1993).

The oleic acid concentration increased from 45.4 to 93% in the top node seed, while oil content and linolenic acid decreased from 14.4 to 26.8% and 5.7 to 34.4% respectively in the top node seed depending on the cultivar. The fully expanded leaves at R5-R6 growth stage showed higher protein and oleic acid concentration, higher nitrate reductase activity, chlorophyll concentration and nitrogen (N) and sulfur (S) percentages in the top node seeds. Seed protein, oil and fatty acids in nodes along the plant depended on the position of node on the main stem, cultivar differences, seed N and S status, and tissue N and S partitioning. Top nodes with higher nitrate reductase activity exhibit high protein and oleic acid (Bellaloui and Gillen, 2010).

Wilcox and Shibles (2001) reported that increase in protein caused decrease in oil and carbohydrates. Of the carbohydrates, sucrose decreased the most with an increase in protein. Temperature during the growing season has an effect on the composition of soybean seeds. Variation in temperature caused changes in soybean oil and carbohydrate contents in controlled growth

chamber studies (Piper and Boote 1999; Thomas *et al.*, 2003) but protein was not affected by temperature changes. Starch and soluble sugar contents decreased with increased temperature and dramatic decreases occurred when the mean air temperature is greater than 20° C (Piper and Boote, 1999; Thomas *et al.*, 2003). Sucrose concentration decreased by 56% with a 15° C temperature increase and stachyose showed a slight reduction while other sugars remained unchanged. Oil content showed positive correlation with temperature (Wolf *et al.*, 1982). Oil content is enhanced with daily average temperature up to approximately 28 °C and decreases with higher temperature in soybean (Piper and Boote, 1999; Thomas *et al.*, 2003).

Water stress influence soybean seed composition by increasing the amount of protein and decreasing the amount of oil in the seed of the severely drought stressed plants (Dornbos and Mullen 1992). Foroud *et al.*, (1993) reported a decrease in protein content of soybean under moisture stress. All cultivars produced smaller seed size with lower protein and higher seed oil, free sugar and sucrose content compared with the same soybean grown in an average year (Poysa and Woodrow, 2002). Sucrose concentration was not affected by water stress in field or greenhouse trials (Egli and Bruening, 2004).

Soybean plant shifts from vegetative to reproductive stages due to changes in length of darkness. Flower initiation in soybean is influenced by temperature. High temperatures during the vegetative period can cause earlier flowering (Elmore and Flowerday, 1984). Late sowing results in reduction in grain yields (Anderson and Vasilas, 1985; Trostle and Bean, 2001). Planting date can influence the composition of soybean seeds. Planting date affects oil content in three Maturity Group II soybean cultivars as revealed by four year study in Wisconsin conducted by Pedersen and Lauer (2003). Early sown crop showed higher oil content at one location while planting date does not affect oil content at other location. Soybean seed protein content is not influenced by planting date (Kane *et al.*, 1997 and Helms *et al.*, 1990). Kentucky and Helms *et al.*, (1990) studied three cultivars in North Dakota and reported that delayed planting increased protein content and reduced oil content.

Planting location can affect the composition of soybean seeds especially protein and oil content (Helms *et al.*, 1998). Effect of location was not significant on sugar content of soybean seeds (Geater and Fehr, 2000) and Cober *et al.*, (1997); however Geater *et al.*, (2000) reported significant effect of location on sugar content. Carrao-Panizzi *et al.*, (1999) reported that the highest isoflavone concentrations were observed in seeds of soybean plants grown in locations with high latitudes (cooler temperatures) when compared to locations with low latitudes (warmer temperatures).

Growth stage and pod position significantly affect the individual and total isoflavone contents (Al-Tawaha, 2010). Isoflavone concentration was lower in seeds collected from the top part of the plants and higher in seeds from the bottom parts (Bordignon *et al.*, 2004). The maximum individual and total isoflavone contents were recorded in soybean seeds from the base parts of plants at brown pod stage (Al-Tawaha, 2010). Nakamura *et al.*, (2001) reported that isoflavone concentration varied with the growth stages of soybean and was highest in mature bean seeds.

Both oil and protein content in soybean seed change due to positional effect (Collins and Cartter, 1956). Protein accumulation was highest in soybean seeds developing at the top parts of plants than the base region while seeds developing at the base parts of plants accumulate greater amount of oil than those from the top (Al-Tawaha, 2010). Incremental trend in protein content was observed in seeds from bottom to top nodes in both normal and high-protein breeding lines (Escalante and Wilcox, 1993a). The glycinin A3 polypeptide content gradually increased in successively lower nodes from the top of the plant while remaining seed storage protein components were not influenced by nodal position (Bennett *et al.*, 2003). Concentration of protein was higher in seeds appearing in the upper one-fourth of the plant while lower oil content than seeds from the lower one-fourth of the plant (Bennett *et al.*, 2003). Variation exists in protein content among the nodes of both determinate and indeterminate plants (Escalante and Wilcox, 1993b). Basal node seeds contain less protein and its concentration increased toward the apical nodes in both types of plants (Escalante and Wilcox, 1993b). Seeds developed on lower nodes showed higher oil in both determinate and indeterminate varieties (Collins and Cartter, 1956). Oil content varies among nodes but plants of the same variety exhibited a similar pattern of oil accumulation. Oleic (18:0) and linoleic (18:2) acids showed difference in accumulation at different nodes (Bennett *et al.*, 2003).

SESAME

A research on Sesame (*Sesamum indicum* L.) was done by Mosjidis and Yermanos, (1985), to study effect of capsule position on seed weight, oil content, and oil composition. Three capsule positions-basal (nodes 16 to 20), middle (nodes 21 to 25), and apical (nodes 26 to 30) were sampled. Middle capsules had heavier seeds and their oil contained more palmitic and oleic acids than seeds from the two sideways capsules. Though, middle capsules had seeds with less oil content than seeds from the side capsules. Seed weight and arachidic acid were found to be stable in seeds from capsules located at different nodes along the plant.

Lipid, protein, RNA and fatty acid composition changes in developing Sesame (*Sesamum indicum*) seeds were determined by Chung *et al.*, (1995). They found that lipid increased from 15 day flowering and continual to 38 DAF (day after flowering) and protein content abruptly increased 12 DAF (day after flowering). RNA increased between 12 to 21 DAF (day after flowering) but after 21 DAF (day after flowering) slowly decreased. Oleic acid, palmitic acid, stearic acid, linoleic acid and linolenic acid consisted 98% of total lipid in growing seed. Oleic acid and linoleic acid are composed of about 80% of total fatty acids. However; Tashiro *et al.*, (1991) found oil content and minor components in the oil of Sesame (*Sesamum indicum*) affected by capsule position; they found seed weight decreased with capsule position from bottom to the top of stem. Oil content was highest at top of main stem and lowest at bottom of main stem.

Sesame seed is a valuable source of oil (44-58%), protein (18-25%), carbohydrate (~13.5%) and ash (~5%) (Borchani *et al.*, 2010). Sesame seed include 50 percent oil with 35% monounsaturated and 44% polyunsaturated fatty acids and 45 percent meal containing 20% protein (Ghandi, 2009; Hansen, 2011). The protein content of sesame seeds ranges from 20 to 30% and is high in tryptophan and methionine (IPGRI, 2004). This unique protein chemistry makes sesame an exceptional protein supplement as compared to soybean, peanuts and other protein sources that are deficient in enough methionine (RMRDC, 2004). Sesame oil is composed of oleic (35.9-47%), linoleic (35.6-47.6), palmitic (8.7-13.8%), stearic (2.1-6.4%), as well as arachidic acids (0.1-0.7%) (Elleuch *et al.*, 2007; Uzun *et al.*, 2002; Weiss, 1983). Sesame seeds include 50-60% oil and nearly 85% unsaturated fatty acids (Latif and Anwar, 2011).

Genetic, climatic and agronomic factors showed immense impact on oil content and composition of sesame. Oil content and composition differs considerably within varieties (Yoshida *et al.*, 2007). Fatty acid composition and oil content are subject to variation by various physiological, ecological and cultural factors (Uzun *et al.*, 2002). Sesame growth rate was negatively influenced by low temperature and less efficient photosynthesis rate (Nath *et al.*, 2003).

The composition of sesame oil varies with the source and depends on factors such as climatic conditions, soil type, maturity of plant and variety (El-Khier *et al.*, 2008). Proximate analysis showed significant variation among the different sesame genotypes which implies that genetic diversity exists among Nigerian sesame (Alege and Mustapha, 2013). Alege *et al.*, (2011) reported significant genetic diversity among three species of sesame from Nigeria using morphological markers. Akinoso

et al., (2006) reported that the oil content varies with genetic and environmental factors.

Sesame, a warm-season annual crop is primarily adapted to areas with long growing season and well-drained soil (Hansen, 2011). It requires long period of sunshine and is generally a short-day plant (Tunde-Akintunde *et al.*, 2012). Some varieties are not influenced by the day-length (Natlund 2002). Sesame requires a constant high temperature. Optimum range for growth, flowering and fruit ripeness is 26-30°C. The minimum temperature for germination is around 12°C, yet even temperatures below 18°C can have a negative effect during germination (Natlund 2002). Pollination and the formation of capsules are inhibited at temperature above 40°C (Tunde-Akintunde *et al.*, 2012).

LINSEED

Linseed (*Linum usitatissimum* L.) is abundant in many nutrients, such as polyunsaturated fatty acid, protein, and lignins (Wang *et al.*, 2007). Linseed contains 36-48 % oil which is the richest source of polyunsaturated fatty acids (PUFA) especially linolenic acid essential in the human diet (Enser *et al.*, 2000; Kouba *et al.*, 2003; Kouba, 2006). Linseed oil is a rich source of unsaturated fatty acids like oleic (C18, 16-24 %), linoleic (C18, 18-24 %) and linolenic acid (C18, 36-50 %) (Flachowsky *et al.*, 1997) and it has a relatively low glucosinolates content (Schuster and Friedt, 1998). Nutritional value of linseed depends upon many factors like cultivar, locality, sowing date and year of production, with cultivar being the most important factor (Oomah *et al.*, 1992). Linseed contains 24.18% crude protein, 37.77% fat, 4.78% crude fiber, 3.50% ash and 25.86% nitrogen free extract, respectively (Khan *et al.*, 2009). The endosperm of linseed contains only 23% of the lipids and 16% of protein (Daun *et al.*, 2003; Oohma, 2003). Flax protein is relatively rich in arginine, aspartic acid and glutamic acid, and the limiting aminoacids are lysine, methionine and cysteine (Chung *et al.*, 2005). Protein content ranges from 18.8 percent to 24.4 percent (Daun and Pryzbylski, 2000).

Temperatures fluctuation before flowering has no effect on composition of the linseed oil. Temperature differential after flowering resulted in drastic change in the fatty-acid composition of the oil. Oil from Dakota variety showed highest linolenic and the lowest oleic acid content at 10°C post-flowering treatment while 26.7°C post-flowering temperature decreased linolenic acid from 57% to 41% and increased oleic acid from 18% to 34% (Yermanos and Goodin, 1965). Both the amount of oil per seed and oil content showed linear decline with daily average temperature between 13 and 25 °C in linseed (Green, 1986).

Rahimi *et al.*, (2011) reported that delayed sowing and increasing nitrogen fertilizer results in decreasing oil content in linseed. Cool temperature and long day length (photoperiod) during and after flowering increase seed yield and linolenic acid. Delayed sowing of linseed reduces number of capsule per plant, number of seeds per capsule, oil content, iodine value and linolenic acid (Ford and Zimmerman, 1964). Longer photoperiods increase oil content, iodine number and linolenic acid while shorter photoperiods result in delayed flowering and lower both, oil content and fatty acids (Sosulski and Gore, 1964). Higher temperatures along with intense radiations favor linseed seed production and oil accumulation. Higher temperature enhances the accumulation of saturated fatty acids while under the condition of low temperature with less intense radiations, plants retain their vegetative stage and yield more fiber of better quality (Rahimi *et al.*, 2011).

Under the warmer temperate climate conditions, linseed exists as a winter plant (Zajac *et al.*, 2012). Cool season and adequate moisture supply during the period from flowering to maturity favor both oil content and quality (Rubilar *et al.*, 2010). With progress in seed formation, level of palmitic and linoleic acid decreases but that of linolenic acid increases in percent while no clear trend has been observed for oleic acid. Environmental conditions required for large grain size also favor higher oil contents and iodine value (Comstock, 1960). Oil content showed linear relationship with polyunsaturated fatty acids namely linoleic and linolenic acids, negative correlation with saturated fatty acids like palmitic and stearic acid and weak correlation with monounsaturated fatty acids (oleic acid) (Adugna and Labuschagne, 2003). Oil content and fatty acid composition showed considerable variation during maturation. Oil content is reported to be increased in first 30 days from flowering. Significant variability in fatty acid levels and total oil content was recorded during 10-20 days after flowering with 3.4% daily increase in oil content. Amount of polyunsaturated fatty acids especially linolenic acid increases while that of palmitic acid decreases during maturation (Bhatia and Sukhija, 1970). Cold weather and lower temperature prevailing during maturation reduce the concentration of palmitic, stearic and oleic acids whereas they enhance the level of linoleic and linolenic acids in 3 normal and 3 genetically modified linseed varieties (Green, 1986).

Day length during growing season showed immense impact on quality of linseed (Nytker *et al.*, 2006). Environmental conditions during the growing season of linseed are major determinant for dry matter accumulation (Marshall *et al.*, 1989; Diepenbrock *et al.*, 1995; Casa *et al.*, 1999; Froment *et al.*, 2000; Hassan *et al.*, 1999; Zajac *et al.*, 2002). Climatic conditions prevailing during two growth

stages i.e. sowing and emergence as well as the plant maturation determine yield (Bravi and Sommovigo, 1997). Yield and its components are influenced by seeding rate of linseed (Zajac *et al.*, 2005). Increasing day temperature from 15 to 27°C and the night temperature from 10 to 22°C results in reduction in oil content to 40 g kg⁻¹ (Berti *et al.*, 2010).

Oil/Fatty acid	Concentration (%) at 15°C	Concentration (%) at 30°C
Linolenic acid	49	31
Palmitic acid	7	37
Oleic acid	28	47
Linoleic acid	15	11
Oil content	38	31
Iodine value	176	140

(Dybing and Zimmerman, 1966)

Linseed can tolerate a wide range of soil types and fertility levels (Townshend and Boleyn, 2013). It requires reasonable soil moisture during development due to its shallow root system and dry weather at harvest (Claridge 1972). Linseed crop can be grown in a variety of environments mainly in temperate climates (Casa *et al.*, 1999; Adugna and Labuschagne, 2003). Temperature during plant development is single determinant influencing seed yield and seed yield components, plant height, time to reach harvest maturity, oil content and oil composition (Dybing *et al.*, 1988; D'Antuono and Rossini, 1995; Casa *et al.*, 1999; Adugna and Labuschagne, 2003; Cross *et al.*, 2003). Seed moisture at harvest may also influence the final seed oil content and the oil composition (Froment *et al.*, 1999; Adugna and Labuschagne, 2003).

Sowing year, location, variety, seeding date, nitrogen fertility and seeding rate are the factors with greatest influence on the final seed yield of linseed (Hassan *et al.*, 1999; Lafond 2001; Adugna and Labuschagne 2002). Nitrogen rate increased oil content and yield. Oil composition was not affected by N, P and K rates, or their interactions. The P and K nutrients did not have an effect on seed yield, oil content, oil yield and oil composition (Berti *et al.*, 2009). Late sown crop encounters higher temperature during reproductive growth resulting in lower seed quality (Greven *et al.*, 2004). Early sowing results in more plant height, number of primary branches, number of capsules/plant, seed/capsule, seed yield, oil % and yield, protein content, higher content of linolenic and linoleic acids (Mirshekari *et al.*, 2012).

Seed traits (Seeds per capsule, thousand seed weight and Seeds-husk ratio) and oil content were influenced by prevailing environmental conditions during maturity and post flowering

period. No. of seeds per capsule, thousand seed weight, seed-husk ratio and oil content was maximum for lowest position and it decreased significantly towards top. Capsule position has significant effect on seed traits and oil contents of linseed that could be due to differential partitioning of photo-assimilate at different points of capsule (Mukhtar *et al.*, 2012). Linseed is a cool temperate annual herb with erect and cylindrical stems. Air temperatures below 10°C possibly inhibit growth rate subsequently delaying the flowering process (Gusta *et al.*, 1997). The growth of capsule or seed depends upon photosynthetic activities of leaves and inflorescences and translocation of photo-assimilate in the plant canopy before flowering (Dordas, 2009; Blum, 1998).

Seed formation and development takes place at different times, different positions (distance from main stem photo-synthates producing region) and at different temperatures, thus physical and chemical characteristics are affected differently. Progressive reduction in number of seeds/ capsule might be the effect of temperature variation. Dry matter accumulation and its partitioning into different plant parts are significantly affected by growing season. Differences in climatic conditions during seed development/maturation and uneven ripening of individual capsules might have caused variations in oil content (Mukhtar *et al.*, 2012). Application of manganese in flax also increased the oil percent and seed yield (Sawan *et al.*, 2001). Khan *et al.* (2011) concluded that in soybean high temperature stress during flowering and pod development had larger effect on seed set and seed filling duration. The comparable study was carried out by Sorensen *et al.*, (2005). They studied changes in fatty acid composition and oil content of two cultivars of Flaxseed (*Linum usitatissimum*). The oil content increased progressively more until about 20 days after flowering (DAF). The total lipid content for both cultivars increased gradually from less than 7% of the dry weight at 8 DAF to over 30% of the dry weight at 20 DAF.

SAFFLOWER

Seed filling pattern of Safflower (*Carthamus tinctorius* L.) affected by seasonal variations and relationship with other agronomic traits were studied by Koutroubas and Papakostab (2010). Plants grown-up under Mediterranean environment subjected to biotic and abiotic stresses throughout the seed filling period reduce their production. Seasonal distinction was pragmatic only for seed filling period that was mostly resolute by the sink size, as it was particular by the number of seeds per head. Seed yield and seed oil yield were considerably connected with seed filling. Likewise, Alizadeh, (2005), studied Safflower germplasm by some agronomic characteristics. Results showed that number of seeds per head and plant height had

recognizable correlation with grain yield, safflower seed-husk ratio decreased from outer to inner side in head. The positional allocation of linolenic acid in triacylglycerols arranged from the immature seeds of safflower 2 days after flowering and from the leaves was unusual. Triacylglycerol was quickly accumulated (14–18 days after flowering). Diacylglycerols and compound lipids reached the highest rate of synthesis 15 days after flowering, and then a maximum incorporation into triacylglycerol occurred 18 days after flowering (Ichihara and Manjiro, 1980). Fatty acid composition of the oil from developing seeds of diverse varieties of safflower was studied by Gecgal *et al.*, (2007). They paid attention on fatty acid composition, moisture and oil content of safflower varieties (Montola-2001 and Centennial). They found that moisture contents turn down 15 days from flowering stage to maturity, as oil content was superior to earlier. The time of the palmitic acid formation decreased in both varieties 15 days from flowering point to maturity, but formation rates of the linoleic and oleic acids greater than before in both varieties.

Zinc and manganese foliar applications significantly increased percentage of palmitic and oleic acids whereas decreased for linoleic acid. Flowering stage is more sensitive to drought than vegetative or grain filling stages. Drought stress during flowering stage imposed the most damage to oil percent and yield and also decreased linoleic acid content and increased stearic and oleic acid content. Foliar application of zinc and manganese can compensate the negative effects of drought on safflower (Dehnaviand Sanavy, 2008). 60 to 70 percent of total oil in safflower seeds is produced at a 22-days period after flowering (Sharma *et al.*, 1995). Drought stress at seed filling stage significantly decreased oil yield and content in sunflower while seed hull percent increased with drought (Mekki *et al.*, 1999). Drought stress at flowering stage resulted in decreasing the unsaturated fatty acids and increasing the saturated fatty acids, while drought stress at seed filling period had no significant effect on seed oil content in safflower. Palmitic acid content was increased and oleic acid content was decreased by drought stress while foliar spray of potassium increased the oleic acid content of oil (Dehnaviand Sanavy, 2008). Application of zinc sulfate increases seed oil and protein content in Indian mustard (Sultana *et al.*, 2001).

Water deficit strikingly decreased the oil percentage, oleic acid and linoleic acid concentrations of the seeds while it increased the protein percentage, palmitic acid and stearic acid in safflower seeds (Mohsennia and Jalilian, 2012). Safflower oil contains the saturated fatty acids: palmitic (C16:0) and stearic acid (C18:0) and the unsaturated fatty acids: oleic (C18:1), linoleic (C18:2) and linolenic acid (C18:3) (Camas and

Esendal, 2006). Safflower seed oil contains about 71–75% linoleic acid, 16–20% oleic acid, 6–8% palmitic acid, and 2–3% stearic acid (Velasco and Fernandez, 2001). The seeds of safflower contain 35–50% oil, 15–20% protein and 35–45% hull fraction (Rahamatalla *et al.*, 2001). Oil quality and yield are both dependent upon the genotype of a plant and its interaction with the environment (Jalilian *et al.*, 2012). Oil contents of Safflower increase significantly with seed development, reach to maximum at 30 DAF and then decrease gradually (Rahmatalla *et al.*, 1998).

Planting date and cultivar showed significant effect on yield and oil content. Delayed sowing decreased yield (Mackinnon and Fettel, 2003). Thin crop density results in lower yield while thick crop and less weeds result in higher yield (Bilgili *et al.*, 2003; Lythoge *et al.*, 2001). Different sowing dates caused flowering and seed development to occur during periods of widely different temperatures, radiation, and day length (Mirshekari *et al.*, 2013). Increased temperature and water stress during seed filling was a major cause of reduction of oil content and thus increased protein content due to late sowing (Hocking and Stapper, 2001). Sowing date can affect oil content and fatty acid composition at the time of seed development (Samanci and O'zkanaynak, 2003). The ratio of oleic and linoleic acids in seed oil is to a large extent dependent upon environmental conditions, particularly moisture and temperature, during seed maturation (Gecgel *et al.*, 2007).

Water deficit stress usually results in shorter plants, less branching, lower seed yield and oil content consequently leading to lower oil yield. Drought stress during seed filling stage in the spring safflower genotypes significantly decreased seed and oil yields in arid and semi-arid regions (Pasban E. 2011). Gecgel *et al.* (2007) revealed that the oil content and the four major fatty acids in safflower seeds were affected by sowing dates, and the availability of water from the flowering to maturity resulted in increased oil content. Ensiye and Khorshid (2010) reported that both oil content and the oil fatty acid composition showed significant differences in relation to the water regime, sowing date and genotype. Sowing date has an expressing influence and determining the appropriate sowing date is one of the most critical factors for optimizing safflower productivity (Yau, 2007, Yarniaet *al.*, 2011).

Conclusion

This review concluded that in oilseed crops, oil content in seed, protein and fatty acid composition of oil is severely affected by environmental factors. Availability of favourable temperature and adequate water are the major factors influencing the quality and quantity of oil. The extent of adverse effect depends upon crop type and its growth stage.

Inadequate nitrogen fixation and higher amounts of precipitation during grain filling stage influence the chemical composition of soybean seeds as increase in oil content and protein content decreases. Temperature fluctuation and drought stress also effects the chemical composition of oil. Linoleic acid and linolenic acid are affected by warmer environments as their content decreases, while the oleic acid concentration increases. Temperature higher than 30°C affects the activity of desaturase enzymes and decreases the sucrose level. Water deficiency causes significant decrease in oil content and increase in protein content. Seed quality is also influenced by pod position on plant. Basal node grains showed lowest level of protein content and high level of oil as compare to apical nodes. Chemical composition of soybean oil is also influenced by its planting location, crop grown at higher latitude has highest level of isoflavone in grains.

In sesame, capsule position significantly affects the concentration and composition of oil. Seeds of middle capsules contain low level of oil contents and high level of palmitic acid and oleic acid in oil than apical and basal capsules. Sesame is long day plant and requires high temperature during all growth stages. Its germination is also significantly affected by temperature below 18°C, while capsule formation and pollination is inhibited above 40°C.

Fatty acid composition of linseed oil is adversely affected by temperature alterations after flowering. Rise in temperature up to 26.7°C decreases the linolenic acid and oleic acid contents. High temperature increases the oil quantity in seed as well as unsaturated fatty acid of oil. Increasing nitrogen level enhances the oil contents but its composition remains unaffected. Oil content, iodine number and linolenic acid concentration are reduced in late sown linseed. Oil composition and concentration is also affected by photoperiod length; thereof longer photoperiod increases the linolenic acid content and vice versa.

Chemical composition of safflower oil is considerably changed due to varying moisture level and temperature. Linoleic acid contents in oil decreases due to moisture stress during flowering, on the other hand stearic acid and oleic acid increases. During seed filling, concentration of palmitic acid increases and oleic acid decreases. Damaging effects of drought can be compensated by foliar application of Zn and Mn. In short, water deficiency decreases the unsaturated fatty acids and increases the saturated fatty acids. Late sowing of crop affect the growing stages of crop specially seed development and flowering stage due to rise in temperature. High temperature during seed development stage increases protein contents and alters the fatty acid composition.

References

- Adugna, W. and M.T. Labuschagne. 2002. Genotype environment interactions and phenotypic stability analyses of linseed in Ethiopia. *Plant Breeding*, 121: 66–71.
- Adugna, W. and M.T. Labuschagne. 2003. Association of linseed characters and its variability in different environments. *J. Agr. Sci.* 140: 285-296.
- Akinoso, R., J.C. Igbeke and T.M.A. Olayanju. 2006. Process optimization of oil expression from Sesame seed (*Sesamum indicum* L.). *The CIGR E. Journal*, 8:1-7.
- Alege, G.O., B.O. Akinyele, S.M. Ayodele and A.V. Ogbode. 2011. Taxonomic Importance of the Vegetative and Pod Characteristics in Nigerian Species of Sesame. *African J. Plant Sci.*, 5 (3): 213-317.
- Alege, G.O. and O.T. Mustapha. 2013. Assessment of genetic diversity in Nigerian sesame using proximate analysis. *G. J. Biosci. Biotechnol.*, 2(1): 57-62.
- Ali, Q., M. Ashraf and F. Anwar F. 2009. Physico-chemical attributes of seed oil from drought stressed sunflower (*Helianthus annuus* L.) plants. *Grasasy Aceites.*, 60 (5), 475-481.
- Alizadeh, K. 2005. Evaluation of Safflower Germplasm by Some Agronomic Characteristics and Their Relationships on Grain Yield Production in the Cold Dry Land of Iran. *Int. J. Agri. Biol.* 7(3): 389–391.
- Al-Tawaha, A.R.M. 2010. Effect of Growth Stage and Pod Position on Soybean Seed Isoflavone Concentration. *Not. Bot. Hort. Agrobot. Cluj.*, 38 (1):92-99.
- Anderson, L.R. and B.L. Vasilas. 1985. Effects of planting date on two soybean cultivars: seasonal dry matter and seed yield. *Crop Sci.*, 25: 999-1004.
- Andrade, F.H. and M.A. Ferreiro. 1996. Reproductive growth of maize, sunflower and soybean at different source levels during grain filling. *Field Crops Res.*, 48: 155–165.
- Anonymous. 2017-18. Economic Survey of Pakistan. Govt. of Pakistan, Finance and Economic Affairs Division, Islamabad, Pakistan.
- Baud, S. and L. Lepiniec. 2010. Physiological and developmental regulation of seed oil production. *Prog Lipid Res.* 49(3): 235-49.
- Baydar, H. and S. Erbas. 2005. Influence of seed development and seed position oil, fatty acids and total tocopherol content in sunflower. *Turkish J. of Agri. Forestry*, 29: 179-186.
- Bellaloui, N. and A.M. Gillen. 2010. Soybean seed protein, oil, fatty acids, N and S partitioning as affected by node position and cultivar differences. *Agric. Sci.* 1:110-118.
- Bellaloui, N., K.N. Reddy, A.M. Gillen, D.K. Fisher and A. Mengist. 2011. Influence of planting date on seed protein, oil, sugars, minerals, and nitrogen metabolism in soybean under irrigated and non-irrigated environments. *Amr. J. Plant Sci.*, 2: 702-715.
- Bennett, J.O., A.H. Krishnan, W.J. Wiebold and H.B. Krishnan. 2003. Positional Effect on Protein and Oil Content and Composition of Soybeans. JF034371L.
- Berti, M., S. Fischer, R. Wilckens, F. Hevia and B. Johnso. 2010. Adaptation and genotype x environment interaction of flaxseed (*Linum usitatissimum* L.) genotypes in south central Chile. *Chilean J. of Agric. Res.*, 70(3): 345-356.
- Berti, M., S. Fischer, R. Wilckens and F. Hevia. 2009. Flaxseed response to N, P, and K fertilization in south central Chile. *Chilean J. Agric. Res.*, 69(2): 145-153.
- Bhattia, I.S. and P.S. Sukhija. 1970. Changes in fatty acids of linseed oil (*Linum usitatissimum* L.) during ripening. *Ind. J. Biochem.*, 7: 215-216.
- Bilgili, U., M. Sincik, A. Uzan and E. Acikgoz. 2003. The influence of row spacing seeding rate on seed yield and yield components of forage turnip. *J. Agron. Crop Sci.*, 189(4): 250-254.
- Blum, A. 1998. Improving wheat grain filling under stress by stem reserve mobilization. *Euphytica*. 100: 77-83.
- Borchani, C., S. Besbes, C.H. Blecker and H. Attia. 2010. Chemical Characteristics and Oxidative Stability of Sesame Seed, Sesame Paste, and Olive Oils. *J. Agric. Sci. Technol.*, 12: 585-596.
- Bordignon, J.R., S.P. Long and N.J. Engeseth. 2004. Isoflavone content in soybean seeds from different parts of plants grown under elevated atmospheric CO₂ or O₃. *Nutraceuticals and Functional Foods: Antioxidants and phytochemical analysis. International Food Safety and Quality Conference, Las Vegas, Nevada.*
- Boydak, E, M. Alpaslan, M. Hayta, S. Gerçek, and M. Simsek. 2002. Seed composition of soybeans grown in the Harran Region of Turkey as affected by row spacing and irrigation. *J. Agric. Food Chem.* 50: 4718-4720.
- Bravi, R.S. and A. Ommovigo. 1997. Seed production and certification of flax/linseed (*Linum usitatissimum* L.). *Sementi-Elette*, 43(2): 5–8.

- Brevedan, R.E. and D.B. Egli. 2003. Crop physiology and metabolism: short period of water stress during seed filling, leaf senescence, and yield of soybean. *Crop Sci.*, 43: 2083-2088.
- Burton, J.W. 1991. Recent Developments in Breeding Soy- Beans for Improved Oil Quality. *Fat Sci. Technol.*, 93: 121-128.
- Camas, N. and E. Esendal. 2006. Estimates of broad-sense heritability for seed yield and yield components of safflower (*Carthamus tinctorius* L.). *Hereditas. J.*, 143: 55-57.
- Carrao-Panizzi, M.C., A.D. Beleia, K. Kitamura and M.C.N. Oliveira. 1999. Effects of genetics and environment on isoflavone content of soybean from different regions of Brazil. *PesqAgropec Bras.* 34: 1787-1795.
- Carrera, C., M.J. Martínez, J. Dardanelli and M. Balzarini. 2009. Water deficit effect on the relationship between temperatures during the seed fill period and soybean seed oil and protein concentrations. *Crop Sci.* 49: 990-998
- Carrera, C., M.J. Martínez, J. Dardanelli, M. Balzarini. 2011. Environmental variation and correlation of seed components in non-transgenic soybeans: protein, oil, unsaturated fatty acids, tocopherols and isoflavones. *Crop Sci.*, 51: 800-809.
- Carver, B.F., J.W. Burton, T.E. Jr-Carter and R.F. Wil- Son. 1986. Response to Environmental Variation of Soybean Lines Selected for Altered Unsaturated Fatty Acid Composition. *Crop Sci.*, 26(6): 1176-1180.
- Casa, R., G. Russell, B. Lo-Cascio and F. Rossini. 1999. Environmental effects on linseed (*Linum usitatissimum* L.) yield and growth of flax at different stand densities. *European J. Agron.*, 11: 267-278.
- Cheesebrough, T.M. 1989. Changes in the enzymes for fatty acid synthesis and desaturation during acclimation of developing soybean seed to altered growth temperature. *Plant Physiol.*, 90: 760-764.
- Chung, H.C., Y.J. Yeea, D.H. Kimb, H.K. Kimb and D.S. Chung. 1995. Changes of lipid, protein, RNA and fatty acid composition in developing sesame (*Sesamum indicum* L.) seeds. *Plant. Sci.*, 109: 237-243.
- Chung, M., B. Lei and E. Li-Chan. 2005. Isolation and structural characterization of the major protein fraction from Norman flaxseed (*Linum usitatissimum* L.). *Food Chem.*, 90: 271-279.
- Claridge, J.H. 1972. Part 6: Flax Chapter 12: Linseed and linen flax. pp. 297-314. In: Arable farm crops of New Zealand. DSIR and A.H & A.W. Reed, Wellington, New Zealand.
- Cober, E.R., J.A. Frégeau-Reid, L.N. Pietrzak, A.R. McElroy, H.D. Voldeng. 1997. Genotype and environmental effects on natto soybean quality traits. *Crop Sci.*, 37: 1151-1154.
- Collins, F.I. and J.L. Cartter. 1956. Variability in chemical composition of seed from different portions of the soybean plant. *Agron. J.*, 48: 216-219.
- Comstock, V.E. 1960. Early generation selection for high oil content and high oil quality in flax. Technical Bulletin 234. University of Minnesota, Agri. Experiment Station. P. 26.
- Cook, D.E. 2008. Genotype-by-Environment Influence on the Traits Associated with Natto Quality. Thesis, Master of Science, Crop and Soil Environmental Sciences, Virginia Polytechnic Institute and State University, Blacksburg, Virginia.
- Cooper, M., D.R. Woodruff, I.G. Phillips, K.E. Basford and A.R. Gilmour. 2001. Genotype-by-management interactions for grain yield and grain protein concentration of wheat. *Field Crops Res.*, 69: 47-67.
- Cross, R.H., S.A.B. McKay, A.G. McHughen and P.C. Bonham-Smith. 2003. Heat-stress effects on reproduction and seed set on *Linum usitatissimum* L. (flax). *Plant Cell and Environ.*, 26: 1013-1020.
- D'Antuono, L.F. and F. Rossini. 1995. Experimental estimation of linseed (*Linum usitatissimum* L.) crop parameters. *Industrial Crops and Products*, 3: 261-271.
- Daun, J., V. Barthet, T. Chornick and S. Duguid. 2003. Structure, composition, and variety development of flaxseed. In: Thompson, L., Cunanne, S. edition. Flaxseed in Human Nutrition. 2nd edition Champaign, Illinois. 1-40.
- Daun, J.K. and R. Przybylski. 2000. Environmental effects on the composition of four Canadian flax cultivars. Proc. 58th Flax Institute, March 23-25, 2000, Fargo, N.D. pp 80-91.
- Dehnavi, M.M. and S.A.M.M. Sanavy. 2008. Effects of withholding irrigation and Foliar Application of Zinc and Manganese on Fatty Acid Composition and Seed Oil Content in Winter Safflower. 7th International Safflower Conference, WaggaWagga, New South Wales, Australia. 1-6.
- Demurin, Y., D. Škori, I. Verešbaranji and S. Joci. 2000. Inheritance of increased oleic acid content in sunflower seed oil. *Helia* 23(32): 87-92.
- Diepenbrock, W., J. Leon and K. Clasen. 1995. Yielding ability and yield stability of linseed in Central Europe. *Agron. J.*, 87: 84-88.

- Dordas, C. 2009. Dry matter, nitrogen and phosphorus accumulation, partitioning and remobilization as affected by N and P fertilization and source - sink relations. *Eur. J. Agron.*, 30(2): 129
- Dornbos, D.L. and R.E. Mullen. 1992. Soybean seed protein and oil contents and fatty acid composition adjustments by drought and temperature. *J. Am. Oil Chem. Soc.*, 69: 228-231.
- Dybing, C.D., P.D. Evenson and C. Lay. 1988. Relationships among daily flower production, length of the flowering period, and seed yield of flax. *Crop Sci.* 28: 287-292.
- Dybing, C.D. and D.C. Zimmerman. 1966. Fatty acid accumulation in maturing flaxseeds as influenced by environment. *Plant Physiol.* 41: 1465-1470.
- Egli, D.B. and W.P. Bruening. 2004. Water stress, photosynthesis, seed sucrose levels and seed growth in soybean. *J. Agric. Sci.*, 142: 1-8.
- El-Khier, M.K.S., K.E.A. Ishag and A.E.A. Yagoub. 2008. Chemical Composition and Oil Characteristics of Sesame Seed Cultivars Grown in Sudan. *Res. J. Agric. Biol. Sci.*, 4(6): 761-766.
- Elleuch, M., S. Besbes, O. Roiseux, C. Blecker, H. Attia. 2007. Quality Characteristics of Sesame Seeds and Byproducts. *Food Chem.*, 103: 641-650.
- Elmore, R.W. and A.D. Flowerday. 1984. Soybean planting date when and why [Online]. NebGuide G84-687-A. Available at <http://ianrpubs.unl.edu/fieldcrops/g687.htm> (verified 2004).
- Enser, M., R.I. Richardson, J.D. Wood, B.P. Gill and P.R. Sheard. 2000. Feeding linseed to increase the n-3 PUFA of pork: Fatty acid composition of muscle, adipose tissue, liver and sausages. *Meat Sci.*, 55: 201-212.
- Ensiye, A. and R. Khorshid. 2010. Effect of Irrigation Regimes on Oil Content and Composition of Safflower (*Carthamus tinctorius* L.) Cultivars. *J. Am. Oil Chem. Soc.*, 87(5): 499-506.
- Escalante, E.E. and J.R. Wilcox. 1993. Variation in seed protein among Nodes of determinate and indeterminate soybean near-isolines. *Crop Sci.*, 33: 1166-1168.
- Escalante, E.E. and J.R. Wilcox. 1993b. Variation in seed protein among nodes of determinate and indeterminate soybean near-isolines. *Crop Sci.*, 33: 1166-1168.
- Escalante, E.E. and J.R. Wilcox. 1993a. Variation in seed protein among nodes of normal- and high-protein soybean genotypes. *Crop Sci.*, 33: 1164-1166.
- Flachowsky, G., T. Langbein, H. Böhme, A. Schneider and K. Aulrich. 1997. Effect of false flax expeller combined with short-term vitamin E supplementation in pigs feeding on the fatty acid pattern, vitamin E concentration and oxidative stability of various tissues. *J. Anim. Physiol. Anim. Nutr.*, 78: 187-195.
- Ford, J.H., D.C. Zimmerman. 1964. Influence of time of flowering on the oil content and oil quality of flaxseed. *Crop Sci.*, 4: 653-656.
- Foroud, N., H.H. Mundel, G. Saindon and T. Entz. 1993. Effect of level and timing of moisture stress on soybean yield, protein, and oil responses. *Field Crops Res.*, 31: 195-209.
- Franklin, P., R. Gardner, B. Pearce and R.L. Mitchell. 2010. Physiology of crop plants. Scientific Press. 336 pp.
- Froment, M.A., J. Smith and K. Freeman. 1999. Influence of environmental and agronomic factors contributing to increased levels of phospholipids in oil from UK linseed (*Linum usitatissimum* L.). *Industrial Crops and Products*, 10: 201-207.
- Froment, M.A., D. Turley and L.V. Collins. 2000. Effect of nutrition on growth and oil quality in linseed. *Test. Agrochem. Cult.*, 21: 29-30.
- Gandhi, A.P. 2009. Simplified process for the production of sesame seed (*Sesamum indicum* L.) butter and its nutritional profile. *Asian J. Food Agro-Indus.*, 2(01): 24-27.
- Geater, C.W. and W.R. Fehr. 2000. Association of total sugar content with other seed traits of diverse soybean cultivars. *Crop Sci.*, 40: 1552-1555.
- Gecgel, U., M. Demirci, E. Esendal, M. Tasan. 2007. Fatty Acid Composition of the Oil from Developing Seeds of Different Varieties of Safflower (*Carthamus tinctorius* L.). *J. Am. Oil Chem. Soc.*, 84: 47-54.
- Ghassemi-Golezani, K. and R. Lotfi. 2013. Influence of Water Stress and Pod Position on Oil and Protein Accumulation in Soybean Grains. *Int. J. Agron. Plant Produc.*, 4 (9): 2341-2345.
- Green, A.G. 1986. Effect of Temperature during Seed Maturation on Oil Composition of Low-Linolenic Genotypes of Flax. *Crop Sci.* 26: 961-965.
- Greven, M.M., B.A. McKenzie, J.G. Hampton, M.J. Hill, J.R. Sedcole and G.D. Hill. 2004. Factors affecting seed quality in dwarf French bean (*Phaseolus vulgaris* L.) before harvest maturity. *Seed Sci. Technol.*, 32: 797-811.

- Grieshop, C.M. and G.C. Fahey. 2001. Comparison of quality characteristics of soybeans from Brazil, China, and the United States. *J. Agri. Food Chem.*, 49: 2669-2673.
- Guleria, S., S. Sucheta, G.S. Balwinder and S.K. Munshi. 2008. Distribution and biochemical composition of large and small seeds of soybean (*Glycine max* L.). *J. Sci. Food Agric.* 88(4): 269-272.
- Gusta, L.V., B.J. OConnor and R.S. Bhatti. 1997. Flax (*Linum usitatissimum* L.) responses to chilling and heat stress on flowering and seed yield. *Can. J. Plant Sci.*, 77: 97-99.
- Hansen, R. 2011. Sesame Profile. 19/08/11. Available at http://www.agmrc.org/commodities_products/grains_oilseeds/sesame_profile.cfm.
- Hassan, F.U., M.H. Leitch and S. Ahmad. 1999. Dry matter partitioning in linseed (*Linum usitatissimum* L.) *J. Agro. Crop Sci.*, 183, 213-216.
- Helms, T.C., T.D. Cai, K.C. Chang and J.W. Enz. 1998. Tofu characteristics influenced by soybean crop year and location [Online]. Available at <http://www.ag.ndsu.nodak.edu/ndagres/fall98/ar31198.htm> (verified 2006).
- Helms, T.C., C.R. Hurburgh, R.L. Lussenden and D.A. Whited. 1990. Economic analysis of increased protein and decreased yield due to delayed planting of soybean. *J. Prod. Agric.* 3: 367-371.
- Hilditch, T.P. and P.N. Williams. 1964. Chemical Constitution of Natural Fats, p.207, Chapman and Hall, London.
- Hobbs, E.H. and H.H. Muendel. 1983. Water requirements of irrigation soybean in southern Alberta. *Can. J. Plant Sci.*, 63: 855-860.
- Hocking, P.J. and M. Stapper. 2001. Effect of Sowing Time and Nitrogen Fertilizer on Canola and Wheat, and Nitrogen Fertilizer on Indian mustard. I. Dry Matter Production, Grain Yield, and yield components. *Aust. J. Agric. Res.*, 52: 623-634.
- Hurburgh Jr, C.R. 1994. Long-term soybean composition patterns and their effect on processing. *J. Am. Oil Chem. Soc.*, 71: 1425-1427.
- Ichihara, K.I. and N. Manjiro. 1980. Fatty acid composition and lipid synthesis in developing safflower seeds. *Phytochem.*, 19 (1): 49-54.
- Illipronti, J.R.R.A., W.J.M. Lommen, C.J. Langerak and P.C. Struik. 2000. Time of pod set and seed position on the plant contribute to variation in quality of seeds within soybean seed lots. *Nether. J. Agric. Sci.*, 48: 165-180.
- International Plant Genetic Resources Institute. 2004. Descriptors for sesame (*Sesamum spp.*) the International Plant Genetic Resources Institute, Rome, Italy: IPGRI.
- Irving, D., M. Shannon, V. Breda and B. Mackey. 1988. Salinity effects on yield and oil quality of high-linoleate and high oleate cultivars of safflower. *J. Agric. Food. Chem.*, 36: 37-42.
- Izquierdo, N.G., G.A.A. Dosio, M. Cantarero, J. Lujan, L.A.N. Aguirrezabal. 2008. Weight per grain, oil concentration, and solar radiation intercepted during grain filling in black hull and striped hull sunflower hybrids. *Crop Sci.*, 48: 688-699.
- Izquierdo, N.G., S. Mascioli, L.A.N. Aguirrezabal and S.M. Nolasco. 2007. Temperature influence during seed filling on tocopherol concentration in a traditional sunflower hybrid. *Grasas y Aceites*. 58: 170-178.
- Kadhem, F.A., J.E. Specht and J.H. Williams. 1985. Soybean irrigation serially timed during stages R1 to R6. II. Yield component responses. *Agro. J.*, 77: 299-304.
- Kane, M.V., C.C. Steele, L.J. Grabau, C.T. MacKown, D.F. Hildebrand. 1997. Early-maturing soybean cropping system: III. Protein and oil contents and oil composition. *Agron. J.*, 89: 464-469.
- Karr-lilienthal, L.K., C.M. Grieshop, J.K. Spears and G.C. Fahey-Jr. 2005. Amino acid, carbohydrate, and fat composition of soybean meals prepared at 55 commercial U.S. soybean processing plants. *J. Agri. Food Chem.*, 53: 2146-2150.
- Khan, M.L., M. Sharif, M. Sarwar and A.M. Sameea. 2009. Chemical Composition of Different Varieties of Linseed. *Pak Vet. J.*, 30(2): 79-82.
- Kouba, M., M. Enser, F.M. Whittington, G.R. Nute and J.D. Wood. 2003. Effect of a high-linolenic acid diet on lipogenic enzyme activities, fatty acid composition, and meat quality in the growing pig. *J. Anim. Sci.*, 81: 1967-1979.
- Kouba, M. 2006. Effect of dietary omega-3 fatty acids on meat quality of pigs and poultry. In: MC Teale (ed). Omega 3 Fatty Acid Research. Nova Publishers. New York, USA: 225-239.
- Koutroubasa, S.D. and D.K. Papakostab. 2010. Seed filling patterns of safflower: Genotypic and seasonal variations and association with other agronomic traits. *Indus. Crops Prod.*, 31(1): 71-76.
- Kumar, V., A. Rani, S. Solanki and S.M. Hussain. 2006. Influence of growing environment on the biochemical composition and physical characteristics of soybean seeds. *J. Food Comp. Anal.*, 19: 188-195.

- Lafond, G. 2001. How is yield determined in flax? *Flax Focus* 14: 6–8. <http://www.flaxcouncil.ca>
- Latif, S. and F. Anwar. 2011. Aqueous enzymatic sesame oil and protein extraction. *Food Chem.*, 125: 679-684.
- Lee, J.D., K.D. Bilyeu and J.G. Shannon. 2007. Genetics and breeding for modified fatty acid profile in soybean seed oil. *J. Crop Sci. Biotech.*, 10(4): 201-210.
- Liu, K.S. 1997. Chemistry and nutritional value of soybean components. Chapter 2. In *Soybeans. Chemistry, Technology and Utilization*. Liu, K., Eds.; Chapman & Hall: New York, NY. 25-113.
- Lobell, S.K. and G.P. Asner. 2003. Climate and management contributions to recent trends in US agricultural yields. *Sci.*, 299: 1032.
- Lythgoe, B., R.M. Norton and D.J. Conner. 2001. Compensatory and competitive ability of two canola cultivars. In: *Proceeding of the Eight Australian Agron. Conference*. 1-8.
- Mackinnon, G.C. and N.A. Fettel. 2003. The effect on sowing time, supplementary water and variety on yield and concentration of canola (*Brassica napus* L.). 13th biennial Australian Research Assembly on Brassicas. *Proceeding of a conference*, Tamworth, New South Wales, Australia. 8-12.
- Maestri, D.M., D.O. Labuckas, J.M. Meriles, A.L. Lamarque, J.A. Zygadlo and C.A. Guzman. 1998. Seed composition of soybean Cultivars evaluated in different environmental regions. *J. Sci. Food Agric.*, 77: 494-498.
- Maleki, A., A. Naderi, R. Naseri, A. Fathi, S. Bahamin and R. Maleki. 2013. Physiological Performance of Soybean Cultivars under Drought Stress. *Bull. Env. Pharmacol. Life Sci.*, 2 (6): 38-44.
- Marshall, G., I.N. Morrison and K. Nawolsky. 1989. Studies on the physiology of (*Linum usitatissimum* L.): The application of mathematical growth analysis: 39– 47. [In:] *Flax: Breeding and utilisation*. Kluwer academic publishers Dordrecht.
- Martin, B.A., R.F. Wilson and R.W. Rinne. 1986. Temperature Effects Upon the Expression of a High Oleic Acid Trait in Soybean. *J. Am. Oil Chem. Soc.*, 63:346-352.
- Matsuzaki, T., S. Iwai and A. Koiwai. 1988. Effects of temperature on seed fatty acid composition in ovary culture of tobacco. *Agric. Biol. Chem.* 52: 1283-1285.
- Mekki, B.B., M.A. EL-Kholy and E.M. Mohamed. 1999. Yield, oil and fatty acids contents as affected by water deficit and potassium fertilization in two sunflower cultivars. *Egyptian J. Agron.*, 21: 67-85.
- Mirshekari, M., R. Amiri, H.I. Nezhad, S.A.S. Noori and O.R. Zanzvaki. 2012. Effect of planting date and low irrigation on quantitative and qualitative traits of flaxseed. *Res. J. Agron.*, 6(1): 20-31.
- Mirshekari, M., N. Majnounhosseini, R. Amiri, A. Moslehi and O.R. Zanzvaki. 2013. Effects of Sowing Date and Irrigation Treatment on Safflower Seed Quality. *J. Agr. Sci. Tech.*, 15: 505-515.
- Mohsennia, O., J. Jalilian. 2012. Response of safflower Seed quality characteristics to different soil fertility systems and irrigation disruption. *Inter. Res. J. Applied Basic Sci.*, 3 (5): 968-976.
- Momen, N.N., R.E. Carlson, R.H. Shaw and O. Arjmand. 1979. Moisture stress effects on the yield components of two soybean cultivars. *Agron. J.*, 71: 86-90.
- Monotti, M. 2003. Growing non-food sunflower in dryland conditions. *Italian J. Agro.* 8: 3-8.
- Mosjidis, J.A. and D.M. Yermanos. 1985. Plant position effect on seed weight, oil content, and oil composition in sesame. *Euphytica*. 34(1): 193-199
- Mukhtar, S., M. Arshad, S.K. Basu, U.H. Fayyaz, M. Ahmed and M. Asif. 2012. Influence of capsule position on seed traits and oil content of linseed (*Linum usitatissimum* L.). *Plant Know. J.* 1(2): 52-56.
- Nakamura, Y., A. Kaihara, K. Yoshii, Y. Tsumura, S. Ishimitsu and Y. Tonogai. 2001. Content and composition of isoflavonoids in mature or immature beans and bean sprouts consumed in Japan. *J. Health Sci.* 47: 394-406.
- Nath, R., P. Chakraborty, P. Bandopadhyay, C. Kundu and A. Chakraborty. 2003. Analysis of relationship between crop growth parameters, yield and physical environment within the crop canopy of sesame (*Sesamum indicum* L.) at different sowing dates, *Arch. Agron. Soil Sci.*, 49: 677-682.
- Naturland. 2002. Organic farming in the Tropics and Subtropics: Sesame. 21/07/11.
- Nian, H., J.L. Wang and Q.K. Yang. 1996. Effects of ecological conditions of the northeast of China on protein, oil and protein, oil content in soybean seeds. *J. Northeast Agric. Univ. (Engl Ed)* 3:1-6.
- Nykter, M., H.R. Kymalainen, F. Gates and A.M. Sjoberg. 2006. Quality characteristics of edible linseed oil. *Agri. Food Sci.*, 15: 402-413.
- Oomah, B.D., G. Mazza, E.O. Kenaschuk. 1992. Cyanogenic compounds in flaxseed. *J. Agric. Food Chem.*, 40: 1346-1348.
- Oomah, B.D. 2003. Processing of flaxseed fiber, oil, protein, and lignan. In: Thompson, L.,

- Cunnane, S. Editores. Flaxseed in Human Nutrition. 2nd. Edn. Champaign, Illinois. 363386.
- Pasban, E.B. 2011. Evaluation of Physiological Indices for Improving Water Deficit Tolerance in Spring Safflower. *J. Agric. Sci. Tech.*, 13: 327-338.
- Pedersen. P., Lauer J.G. 2003. Soybean agronomic response to management systems in the upper midwest. *Agron. J.*, 95: 1146–1151.
- Pedersen. P. and J.G. Lauer. 2004. Response of soybean yield components to management system and planting date. *Agron. J.* 96: 1372–1381.
- Piper, E.L. and K.J. Boote. 1999. Temperature and cultivar effects on soybean seed oil and protein concentrations. *J. Amer. Oil Chem. Soc.* 76: 1233-1241.
- Poysa, V. and L. Woodrow. 2002. Stability of soybean seed composition and its effect on soymilk and tofu yield and quality. *Food Res. Int.* 35: 337-345.
- Pritchard, F.M., H.A. Eagles, R.M. Norton, P.A. Salisbury and M. Nicolas. 2000. Environmental effects on seed composition of Victorian canola. *Aust. J. Exp. Agric.*, 40(5): 679-685.
- Pritchard, F.M., R.M. Norton, H.A. Eagles, P.A. Salisbury and M. Nicolas. 1999. The effect of environment on victorian canola quality. 10th International rapeseed congress, Caneberra, Australia.
- Qadir, G., S. Ahmad, U.H. Fayyaz and M.A. Cheema. 2006. Oil and fatty acid accumulation in sunflower as influenced by temperature variation. *Pak. J. Bot.*, 38(4): 1137-1147.
- Rahamatalla, A.B., E.E. Babiker, A.G. Krishna and A.H. El-Tinay. 1998. Changes in chemical composition, minerals and amino acids during seed growth and development of four safflower cultivars. *Plant Food Hum. Nutr.*, 52: 161-170.
- Rahamatalla, A.B., E.E. Babiker, A.G. Krishna and A.H. El-Tinay. 2001. Changes in fatty acids composition during seed growth and physicochemical characteristics of oil extracted from four safflower cultivars. *Plant Foods Hum. Nutr. J.*, 56: 385–395.
- Rahimi, M.M., G. Nourmohammadi, A. Ayneband, E. Afshar and G. Moafpourian. 2011. Study on planting date and nitrogen levels on yield, yield components and fatty acid of linseed (*Linum usitatissimum* L.). *World Appl. Sci. J.*, 12(1): 59-67.
- Ram Materials Research and Development Council. 2004. Survey report of ten selected agro raw materials in Nigeria, Raw Materials Research and Development Council, Abuja, Nigeria, 89.
- Roche, J., A. Bouniols, Z. Mouloungui, T. Barranco and M. Cerny. 2006. Management of environmental crop conditions to produce useful sunflower oil components. *Eur. J. Lipid Sci. Technol.*, 108: 287–297.
- Rose, I.A. 1988. Effects of moisture stress on the oil and protein components of soybean seeds. *Aust. J. Agri. Res.* 39: 163-170.
- Rubilar, M., C. Gutiérrez, M. Verdugo, C. Shene and J. Sineiro. 2010. Flaxseed as a source of functional ingredients. *J. Soil Sci. Plant Nutr.*, 10 (3): 373-377.
- Samanci, B. and Q. Ozkaynak. 2003. Effect of Planting Date on seed yield, oil content and fatty acid composition of safflower (*Carthamus tinctorius* L.) cultivars grown in Mediterranean region of turkey. *J. Agron. Crop Sci.*, 189(5): 359-366.
- Santalla, G., I. Riccobene, L. Aguirrezábal and S. Nolasco. 1995. Influencia de la radiación solar interceptada durante el llenado de los frutos. 3. Variaciones de la composición acidica dentro del capítulo. 1 ° Congreso Nacional de Soja y 2 ° Congreso Nacional de Oleaginosos, Pergamino, Argentina 2: 15 –22.
- Sawan, Z.M., S.A. Hafez and A.E. Basyony. 2001. Effect of phosphorus fertilization and foliar application of chelated zinc and calcium on seed, protein and oil yields and oil properties of cotton. *J. Agric. Sci.*, 136(2): 191-198.
- Schmitt, M.A., J.A. Lamb, R.W. Gyles, J.H. Orf and G.W. Hehm. 2001. In-season fertilizer nitrogen applications for soybean in Minnesota. *Agric. J.*, 93: 983-988.
- Schuster, A. and W. Friedt. 1998. Glucosinolate content and composition as parameters of quality of *Camelina* seed. *Ind. Crops Prod.*, 7: 297– 302.
- Schuster, W. and J. Boehm. 1981. Experience in soybean breeding in Middle Europe, in Production and Utilization of Protein in Oilseed Crops. Ed by Bunting ES, Martin Nijhoff, The Hague, 158-171.
- Sexton, P.J., N.C. Paek and R. Shibles. 1998. Soybean sulfur and nitrogen balance under varying levels of available sulfur. *Crop Sci.*, 38: 975-982.
- Shafii, B., K.A. Mahler, W.J. Price and D.L. Auld. 1992. Genotype x environment interaction effects on winter rapeseed yield and oil content. *Crop Sci.*, 32: 922-927.
- Sharma, C.P., K. Neena, C. Chatterjee and N. Kurana. 1995. Manganese stress change physiology and oil content of linseed (*Linum usitatissimum* L.). *Indian J. Exp. Biol.*, 3: 701-704.

- Simolenka, G. and P.J.C. Kuiper. 1977. Effect of low temperature upon lipid and fatty acid composition of roots and leaves of winter rape plants. *Plant Physiol.*, 41: 29-35.
- Singh, S.P., K.N. Nansal and V. Nepalia. 2001. Effect of nitrogen, its application time and sulphur on yield and quality of soybean (*Glycine max*L.). *Ind. J. Agron.*, 46: 141-144.
- Sosulski, F.W. and R.F. Gore. 1964. The effect of photoperiod and temperature on the characteristics of flaxseed oil. *Can. J. Plant Sci.*, 44: 381-382.
- Steer, B. and G. Seiler. 1990. Changes in fatty acid composition of sunflower (*Helianthus annuus*L.) seeds in response to time of nitrogen application, supply rates and defoliation. *J. Sci. Food Agric.*, 51: 11-26.
- Strecker, L., M. Bieber, A. Maza, T. Grossberger and W. Doscoczynski. 1997. Aceite de maíz. Antecedentes, composición, procesamiento, refinación, utilización y aspectos nutricionales. *Aceites y Grasas*, 507-527.
- Sultana, N., T. Ikeda and M.A. Kashem. 2001. Effect of foliar spray of nutrient solutions on photosynthesis, dry matter accumulation and yield in sea water-stressed rice. *Environmental and Experimental Botany*, 46: 129-140.
- Sweeney, D.W. and G.V. Granade. 1993. Yield, nutrient, and soil sulfur response to ammonium sulfate fertilization of soybean cultivars. *J. Plant Nutr.*, 16: 1083-1098.
- Tang, G.O., W.P. Novitzky, H.C. Griffin, S.C. Huber and R.E. Dewey. 2005. Oleate Desaturase Enzymes of Soybean: Evidence of Regulation through Differential Stability and Phosphorylation. *Plant J.* 44(3): 433-446.
- Tashiro, T., F. Yasuko and O. Tashihiko. 1991. Oil content of seeds and minor components in the oil of Sesame (*Sesamum indicum* L.). As affected by capsule position. *Japan J. Crop Sci.*, 60(1): 116-121.
- Thomas, J.M.G., K.J. Boote, L.H. Allen-Jr, M. Gallo-Meagher and J.M. Davis. 2003. Elevated temperature and carbon dioxide effects on soybean seed composition and transcript abundance. *Crop Sci.* 43: 1548-1557.
- Thompson, J.A. 1978. Effect of irrigation interval and plant population on growth, yield and water stress of soybean in semi-arid environmental. *Aust. J. Exp. Agric. Husbandry*, 18: 276-281.
- Thuzar, M., A.B. Puteh, N.A.P. Abdullah, M.B. Mohd and K.J. Lassim. 2010. The effects of temperature stress on the quality and yield of soya bean (*Glycine max* L.) Merrill. *J. Agri. Sci.*, 2: 172-179.
- Townshend, J.M. and J.M. Boleyn. 2013. Evaluation of two new linseed cultivars for oilseed production in Canterbury. Agronomy Society of New Zealand Special Publication No. 13 / Grassland Research and Practice Series No. 14. pp. 105-112. Available on http://www.grassland.org.nz/publications/nzgrassland_publication_2495.pdf.
- Tremolieres, H., A. Tremolieres and P. Mazliak. 1978. Effects of light and temperature on fatty acid desaturation during the maturation of rapeseed. *Phytochem.* 17: 685-687.
- Trostle, C. and B. Bean. 2001. Effect of planting date and maturity group on soybean yield in the Texas south plains in 2001. [Online]. Texas Cooperative Extension. Available at <http://lubbock.tamu.edu/othercrops/soybeandat.htm> (verified 6 Nov 2006)
- Tunde-Akintunde, T.Y., M.O. Oke and B.O. Akintund. 2012. Sesame Seed. In: Oilseeds. Ed. U.G. Akpan. Janeza Trdine 9, 51000 Rijeka, Croatia. pp. 81-98.
- Uzun, B., S. Ulger and M.I. agirgan. 2002. Comparison of Determinate and Indeterminate Types of Sesame for Oil Content and Fatty Acid Composition. *Turk J. Agric. For.*, 26: 269-274.
- Velasco, L. and J.M. Fernandez-Martinez. 2001. Breeding for oil quality in safflower. (ed. Bergman JW, Mündel HH), pp 133-137. Proceedings of the 5th International Safflower Conference. Williston, North Dakota and Sidney, Montana, USA.
- Vollmann, J., C.N. Fritz, H. Wagentristl and P. Ruckebauer. 2000. Environmental and genetic variation of soybean seed protein content under Central European growing conditions. *J. Sci. Food Agric.*, 80: 1300-1306.
- Wang, B., D. Li, L.J. Wang, Z. Huang, L. Zhang, X.D. Chen and Z. Mao. 2007. Effect of Moisture Content on the Physical Properties of Fibered Flaxseed. *Inter. J. Food Eng.*, 3(5): 1-1.
- Weiss, E.A. 1983. Sesame. In: "Oilseed Crops". Longman Inc., New York. 282340.
- Wesley, T.L., R.E. Lamond, V.L. Martin and S.R. Duncan. 1998. Effects of late-season nitrogen fertilizer on irrigated soybean yield and composition. *J. Prod. Agric.*, 11: 331-336.
- Wilcox, J.R. and J.F. Cavins. 1992. Normal and low linolenic acid soybean strains: response to planting date. *Crop Sci.* 32: 1248-1251.

- Wilcox, J.R. and R.M. Shibles. 2001. Interrelationships among seed quality attributes in soybean. *Crop Sci.*, 41:11-14.
- Wilson, J.M. and R.M.M. Crawford. 1974. Leaf fatty acid content in relation to hardening and chilling injury. *J. Exp. Bot.*, 25: 121-131.
- Wilson, R. 2004. Seed composition. In: STEWART, B.A.; NIELSEN, D.R. (Ed.). Soybeans: improvement, production and uses. 3rd ed. Madison: ASA/CSSA/SSSA. 621-677. (Agronomy monograph, 16).
- Wolf, R.B., J.F. Canvins, R. Kleiman and L.T. Black. 1982. Effect of Temperature on soybean seed constituents: oil, protein, moisture, fatty acids, amino acid and sugars. *J. Am. Oil Chem. Soc.*, 59: 230-232.
- Yarnia, M., M.B. Khorshidi, E. Benam, M.T. Farajzadeh, N. Nobari and V. Ahmadzadeh. 2011. Effect of planting dates and density in drought stress condition on yield and yield components of Amaranth cv. Koniz. *Adv. Envi. Biol.* 5(6): 1139-1149.
- Yau, S.K. 2007. Winter versus spring sowing of rain-fed safflower in a semi-arid, high-elevation Mediterranean environment. *Euro. J. Agron.*, 26: 249-256.
- Yermanos, D.M. and J.R. Gooden. 1965. Effect of Temperature during Plant Development on the Fatty Acid Composition of Linseed Oil. *Agron. J.*, 57: 453-454.
- Yoshida, H., M. Tanaka, Y. Tomiyama and Y. Mizushima. 2007. Regional distribution in the fatty acids of triacylglycerols and phospholipids of sesame seeds (*Sesamum indicum* L.). *J. Food Lipids.*, 14: 189-201.
- Zajac, T., S. Grzesiak, B. Kulig and M. Polacek. 2005. The estimation of productivity and yield of linseed (*Linum usitatissimum* L.) using the growth analysis. *Acta Physiol. Plant.*, 27(4A): 549-558.
- Zajac, T., K. Klima, F. Borowiec, R. Witkowicz and J. Barteczko. 2002. Yielding of linseed varieties in various site conditions. *Rosliny Oleiste XXIII*: 275-286. (In Polish).
- Zajac, T., A. Oleksy, A. Klimek-Kopyra and B. Kulig. 2012. Biological determinants of plant and crop productivity of flax (*Linum usitatissimum* L.). *Acta. Agrobotanica.*, 65(4): 3-14.